

Math 111.01 Summer 2003  
July 16, 2003

The goal of this problem is to have you prove the chain rule.

Recall that the definition of derivative is  $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ . If  $h$  is small, then we can approximate  $f'(x)$  by  $\frac{f(x+h) - f(x)}{h}$ .

Suppose that we know  $f(x)$  and  $f'(x)$ . We can use these to approximate the value of  $f(x+h)$  for small  $h$ .

Solving  $f'(x) \approx \frac{f(x+h) - f(x)}{h}$  for  $f(x+h)$  yields  $f(x+h) \approx f(x) + hf'(x)$ . This is called a *linear approximation*.

**Example:** Suppose that we want to approximate the value of  $(7.1)^2$ . We know that  $7^2 = 49$ . We can use the idea above by letting  $f(x) = x^2$ . Then  $f'(x) = 2x$ . Use the formula above with  $h = 0.1$  and  $x = 7$ . Thus,  $f(x+h) \approx f(x) + hf'(x)$  or  $(x+h)^2 \approx x^2 + 2hx$ .

Therefore,  $(7 + .1)^2 \approx 7^2 + 2(.1)(7) = 49 + 1.4 = 50.4$ .

**Use a linear approximation to prove the chain rule. That is, prove that  $[f(g(x))]' = f'(g(x)) \cdot g'(x)$ .**

*(Hint: Differentiate  $f(g(x))$  using  $g(x+h) \approx g(x) + hg'(x)$  and  $f(z+k) \approx f(z) + kf'(z)$ .)*