

Solutions should be completed, but not submitted, by Wednesday, September 30, 2015.

1. The purpose of this exercise is to lead you through an alternate verification that the density of a $\beta(a, b)$ random variable, namely

$$f(x) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} x^{a-1}(1-x)^{b-1}, \quad 0 < x < 1,$$

is, in fact, a legitimate density. Suppose that

$$B(a, b) = \int_0^1 x^{a-1}(1-x)^{b-1} dx.$$

Our goal is to show that

$$B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}.$$

(a) As a first step, we will need to prove that

$$B(a, b) = 2 \int_0^{\pi/2} \cos^{2a-1}(\theta) \sin^{2b-1}(\theta) d\theta.$$

This is done by making the substitution $x = \cos^2(\theta)$.

(b) Now consider

$$\Gamma(a)\Gamma(b) = \int_0^\infty u^{a-1}e^{-u} du \cdot \int_0^\infty v^{b-1}e^{-v} dv = \int_0^\infty \int_0^\infty u^{a-1}v^{b-1}e^{-(u+v)} du dv.$$

Change variables by letting $u = x^2$ and $v = y^2$ to show that

$$\Gamma(a)\Gamma(b) = 4 \int_0^\infty \int_0^\infty x^{2a-1}y^{2b-1}e^{-(x^2+y^2)} dx dy.$$

(c) Change to polar coordinates with $x = r \cos(\theta)$, $y = r \sin(\theta)$ for $0 \leq r < \infty$, $0 \leq \theta \leq \pi/2$ to show that

$$\Gamma(a)\Gamma(b) = 4 \int_0^\infty r^{2a+2b-2}e^{-r^2} r dr \int_0^{\pi/2} \cos^{2a-1}(\theta) \sin^{2b-1}(\theta) d\theta. \quad (*)$$

(d) Let $t = r^2$ to show that

$$2 \int_0^\infty r^{2a+2b-2}e^{-r^2} r dr = \int_0^\infty t^{a+b-1}e^{-t} dt = \Gamma(a+b).$$

(e) Combine (a) and (d) to conclude from (*) that

$$\Gamma(a)\Gamma(b) = \Gamma(a+b)B(a, b)$$

as required.